JOURNAL
OF THE AIR & WASTE MANAGEMENT ASSOCIATION VOLUME 47

Subject Index

Accuracy Assessments
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976

Acid Deposition
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Preceptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

Acidity
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411

Aeration Units
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Aerosols
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
Spatial and temporal patterns in particle data measured during the MOHAVE Study, 119
Analyses of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
Methods for real-time, in situ measurement of aerosol light absorption, 157
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Formulation of a second-generation relative plume visibility model, 176
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238
São Paulo aerosol characterization study, 1297

Agricultural Soils
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

Air Bags
Henry's law coefficient of hydrazoic acid, 1216

Air & Waste Management Association
Air quality monitoring during construction and initial occupation of a new building, 890

Air Monitoring
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: summary of data and practical guidelines, 20

Air Pollution
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Modeling anthropogenic trends in air quality data, 66
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Preceptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524
An analysis of Michigan and California CO remote sensing measurements, 601
Redundant measurements of urban air monitoring networks in air quality reporting, 614
Comparing neural networks and regression models for ozone forecasting, 653

Kinetic behavior of the toluene biofiltration process, 664
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Status and future of baghouses in the utility industry, 704
Formation and transport of the Madrid ozone plume, 766
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871
Water content dynamics in biofiltration: The role of humidity and microbial heat generation, 936
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Transient behavior of the biofilters for toluene removal, 1142
A new method for measuring vapor emissions, 1161
Integrated environmental control modeling of coal-fired power systems, 1180
Application of computational methods for the determination of traffic emissions, 1204
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291
São Paulo aerosol characterization study, 1297

Air Pollution Standards
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468
Attainment flip-flops and the achievability of the ozone air quality standard, 620

Air Quality
Preceptor model applied to patterns in space (RMAPS) Part I-Model description, 216
Preceptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Redundant measurements of urban air monitoring networks !n air quality reporting, 614
Comparing neural networks and regression models for ozone forecasting, 653
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
São Paulo aerosol characterization study, 1297

Air Quality Assessment
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302

Air Quality Standards
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468
Attainment flip-flops and the achievability of the ozone air quality standard, 620
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070
Health risk assessment of fluctuating concentrations using lognormal models, 1152

Air Quality Studies
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
The effect of California's wintertime oxygenated fuels program on ambient carbon monoxide concentrations, 775
Air quality monitoring during construction and initial occupation of a new building, 890
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

Air Quality Trends Report
Modeling anthropogenic trends in air quality data, 66
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

Air Toxics
Evaluating design options for biofilters, 37
Formulation of a second-generation relative plume visibility model, 176
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
Airborne hexavalent chromium in Southwestern Ontario, 905
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Alternative/Innovative Technologies
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
A new method for measuring vapor emissions, 1161
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Alternative Transportation Fuels
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Alveolar Breath
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Alternative/Innovative Technologies
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Ambient Air
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
Comparison of selenium and sulfur at remote sites throughout the United States, 204
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Measurement of ozone concentrations in ambient air using a badge-type passive monitor, 697
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Ambient Air Quality Standards
Redundant measurements of urban air monitoring networks in air quality reporting, 614
Attainment flip-flops and the achievability of the ozone air quality standard, 620
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Ambient Concentrations
Spatial and temporal patterns in particle data measured during the MOHAVE Study, 119
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
Comparison of selenium and sulfur at remote sites throughout the United States, 204
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
The effect of California's wintertime oxygenated fuels program on ambient carbon monoxide concentrations, 775
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238
São Paulo aerosol characterization study, 1297

Ammonia
Bioconversion of methylamine in biofilters, 58

Ammonium
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Analytical Methods
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Area Sources
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Arsenic
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Partitioning of arsenic species in fine-grained soils, 481

Ash, Bottom
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Ash, Fly
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Atmospheric Chemistry
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Formulation of a second-generation relative plume visibility model, 176
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Airborne hexavalent chromium in Southwestern Ontario, 905
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Henry's law coefficient of hydrazoic acid, 1216

Atmospheric Transport
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Automotive Exhaust
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
An analysis of Michigan and California CO remote sensing measurements, 601
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
Application of computational methods for the determination of traffic emissions, 1204

Azides
Henry's law coefficient of hydrazoic acid, 1216

Back-Trajectories
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Background Air
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592

Baghouses
Status and future of baghouses in the utility industry, 704

Batch Biofiltration
Water content dynamics in biofiltration: The role of humidity and microbial heat generation, 936

Better Utilization
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Biofilter
Bi conversion of methylamine in biofilters, 58

Biofiltration
Evaluating design options for biofilters, 37
Kinetic behavior of the toluene biofiltration process, 664
Screening of biofiltering material for VOC treatment, 674
Transient behavior of the biofilters for toluene removal, 1142

Biomarkers
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Bioremediation
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710

Biotransformation
Biotransformation of herbicides in the presence of various electron acceptors, 690

Blood
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Blood/Breath Partition Coefficient
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Body Burden
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

BOF Dust
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Bootstrap Methods
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Calculation Methods
Comparing neural networks and regression models for ozone forecasting, 653
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Application of computational methods for the determination of traffic emissions, 1204

Calibration
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Canada
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Canopy Resistances Estimates
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in Southern Europe, 792

Carbon Dioxide
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806
Air quality monitoring during construction and initial occupation of a new building, 890

Carbon Monoxide
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
An analysis of Michigan and California CO remote sensing measurements, 601
The effect of California's wintertime oxygenated fuels program on ambient carbon monoxide concentrations, 775
Air quality monitoring during construction and initial occupation of a new building, 890
Characterization of air quality problems in five finnish indoor ice arenas, 1979
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

Carbonaceous Material
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Cement Kilns
Destruction of chlorofluorocarbons in a cement kiln, 1220

CERCLA
On the Impact of Future Land Use Assumptions on Risk Analysis for Superfund Sites, 881
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Chamber
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Characterization
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Combustion and gasification properties of plastics particles, 861
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Chemical Mass Balances
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Chlorofluorocarbons
Destruction of chlorofluorocarbons in a cement kiln, 1220

Clean Air Act Amendments
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468

Clean Air Corridors
Clean air corridors: A geographic and meteorologic characterization, 403

Cleanup Standards
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Climate Change
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806

Coal-Burning Facilities
Integrated environmental control modeling of coal-fired power systems, 1180

Coal Tar Pitch
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Combustion
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Combustion and gasification properties of plastics particles, 861
Simultaneous treatment of industrial wastewater and vent streams, 961
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Compartmental Modeling
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Compliance
Redundant measurements of urban air monitoring networks in air quality reporting, 614

Composting
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

Confidence Intervals
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976

Consumer Products
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Contaminated Soil
Partitioning of arsenic species in fine-grained soils, 481
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Continuous Emission Monitoring
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Control Technologies
Kinetic behavior of the toluene biofiltration process, 664
Status and future of baghouses in the utility industry, 704
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Transient behavior of the biofilters for toluene removal, 1142
Integrated environmental control modeling of coal-fired power systems, 1180

Control Strategies
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Integrated environmental control modeling of coal-fired power systems, 1180
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Costs
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Integrated environmental control modeling of coal-fired power systems, 1180

CT
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Data Analysis
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Data Collection
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Data Quality
Analyses of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Decision Making
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Integrated environmental control modeling of coal-fired power systems, 1180
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Desert Southwest
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Destruction and Removal Efficiencies
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976
Destruction of chlorofluorocarbons in a cement kiln, 1220
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Diffusion Denuder Samplers
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

JOURNAL
OF THE AIR & WASTE MANAGEMENT ASSOCIATION
VOLUME 47

Dimensional Analysis
Simultaneous treatment of industrial wastewater and vent streams, 961

Dispersion Models
Formulation of a second-generation relative plume visibility model, 176
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Destruction and Removal Efficiencies
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Dust, Fugitive
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Economics
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean air act amendments, 395
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Electric Utilities
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Status and future of baghouses in the utility industry, 704
Integrated environmental control modeling of coal-fired power systems, 1180

Electron Beam
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Electrostatic Precipitation
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Emission Controls
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
An analysis of Michigan and California CO remote sensing measurements, 601
Kinetic behavior of the toluene biofiltration process, 664
Transient behavior of the biofilters for toluene removal, 1142
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Emission Inventories
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
Application of computational methods for the determination of traffic emissions, 1204

Emission Rates
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
Exhaust emissions from four-stroke lawn mower engines, 945
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111
A new method for measuring vapor emissions, 1161

Emissions Reduction
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Emission Standards
An analysis of Michigan and California CO remote sensing measurements, 601
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Emission Trends
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

Emissions
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871
Exhaust emissions from four-stroke lawn mower engines, 945
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Emissions, Automotive
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
Application of computational methods for the determination of traffic emissions, 1204
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Emissions, Carbon Monoxide
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Emissions Characterization
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Emissions, Hydrocarbon
A new method for measuring vapor emissions, 1161
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Emissions Inventory
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Emissions Management
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302

Emissions, Automotive
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
An analysis of Michigan and California CO remote sensing measurements, 601
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Emissions, Carbon Monoxide
An analysis of Michigan and California CO remote sensing measurements, 601

Emissions, Fugitive
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Emissions, Hydrocarbon
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Emissions Reduction
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Environmental Management
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Comparing neural networks and regression models for ozone forecasting, 653
Combustion and Gasification Properties of Plastics Particles, 861
On the impact of future land use assumptions on risk analysis for superfund sites, 881
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Health risk assessment of fluctuating concentrations using lognormal models, 1152

Epidemiology
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Ethyl Corp.
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Ethylene-Vinyl Acetate Copolymer (EVA)
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Exposure
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Exposure Parameters
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
On the impact of future land use assumptions on risk analysis for superfund sites, 881
Air quality monitoring during construction and initial occupation of a new building, 890
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Small sample properties of nonparametric bootstrap t confidence intervals, 1197
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Fabric Filters
Status and future of baghouses in the utility industry, 704

Facility Siting
Redundant measurements of urban air monitoring networks in air quality reporting, 614

Fault-Mode Incineration
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953

Fenton's Reagent
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710

Fine Particle Carbonaceous Material
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Flue Gases
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Fluidized-Bed Incinerator
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953

Fluoride
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Fly Ash
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Forests
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

FTIR
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Fuels
Combustion and Gasification Properties of Plastics Particles, 861

Fuels, Alternative
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Combustion and Gasification Properties of Plastics Particles, 861

Fuels, Fossil
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212

Gangue Constituents
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Gas Chromatography
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Gas Chromatography-Mass Spectrometry
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Gases, Greenhouse
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806

Gas, Natural
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

Gasoline
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

Geography
Comparing neural networks and regression models for ozone forecasting, 653

Global Warming/Greenhouse Effect
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Halocarbons
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Hazardous Air Pollutants
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468
Air quality monitoring during construction and initial occupation of a new building, 890
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111
Transient behavior of the biofilters for toluene removal, 1142
Henry's law coefficient of hydrazoic acid, 1216
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Hazardous Waste
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Hazardous Waste Cleanup
Partitioning of arsenic species in fine-grained soils, 481
Biotransformation of herbicides in the presence of various electron acceptors, 690
On the impact of future land use assumptions on risk analysis for superfund sites, 881
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Hazardous Waste Management
On the impact of future land use assumptions on risk analysis for superfund sites, 881
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Hazardous Waste Sites
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Haze
Spatial and temporal patterns in particle data measured during the MOHAVE Study, 119
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
Methods for real-time, in situ measurement of aerosol light absorption, 157
Formulation of a second-generation relative plume visibility model, 176
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317

Health Risk Assessment
Health risk assessment of fluctuating concentrations using lognormal models, 1152

Health Effects/Risks
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
On the Impact of Future Land Use Assumptions on Risk Analysis for Superfund Sites, 881
Airborne hexavalent chromium in Southwestern Ontario, 905
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Health risk assessment of fluctuating concentrations using lognormal models, 1152
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Henry's Law
Henry's law coefficient of hydrazoic acid, 1216

Hexavalent Chromium
Background air concentrations of Cr(VI) in Hudson County, New

Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592

High Solids Degradation
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

HON Rule
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468

Human Exposure
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
Air Quality Monitoring During Construction and Initial Occupation of a New Building, 890
Airborne hexavalent chromium in Southwestern Ontario, 905
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095
Henry's law coefficient of hydrazoic acid, 1216
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Humidity
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
Water content dynamics in biofiltration: The role of humidity and microbial heat generation, 936

Hydrocarbons
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Ice Arena
Characterization of air quality problems in five finnish indoor ice arenas, 1979
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

IMPROVE
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Incineration
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Combustion and Gasification Properties of Plastics Particles, 861
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Individual Exposures
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Indoor Air Pollution
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Airborne hexavalent chromium in Southwestern Ontario, 905
Characterization of air quality problems in five finnish indoor ice arenas, 1079
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Industrial Pollution
Henry's law coefficient of hydrazoic acid, 1216

Inhalation
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Innovative Technologies
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
Integrated environmental control modeling of coal-fired power systems, 1180
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Inspection/Maintenance
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426

Interlaboratory Study
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Landfills
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Langmuit Adsorption
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953

Leaching
Partitioning of arsenic species in fine-grained soils, 481
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Leaded Gasoline
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Light Absorption
Methods for real-time, in situ measurement of aerosol light absorption, 157

Light Extinction
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317

Limestone Scrubbing
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Long Range Transport
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331

Lower Fraser Valley
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley

Major Phases
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Manganese
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Mathematical Modelling
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Measurement Techniques
Methods for real-time, in situ measurement of aerosol light absorption, 157
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Airborne hexavalent chromium in Southwestern Ontario, 905
A new method for measuring vapor emissions, 1161
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Media Screening
Screening of biofiltering material for VOC treatment, 674

Mercury
Formulation of a second-generation relative plume visibility model, 176

Mesoscale Transport
Summer transport patterns affecting the Mohave power project emissions, 383

Meteorological Conditions
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269
Summer transport patterns affecting the Mohave power project emissions, 383
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419
Comparing neural networks and regression models for ozone forecasting, 653

Methyl Ethyl Ketone
Screening of biofiltering material for VOC treatment, 674

Methyl Teritary Butyl Ether (MTBE)
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Methylamine
Bioconversion of methylamine in biofilters, 58

Metropolitan Areas
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Comparing neural networks and regression models for ozone forecasting, 653
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

MGP-REM Process

Microanalysis
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Microtox
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Mixing Heights
Summer transport patterns affecting the Mohave power project emissions, 383

MMT
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Mobile Sources
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
An analysis of Michigan and California CO remote sensing measurements, 601
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
Effects of Grades and Other Loads on On-Road Emissions of Hydrocarbons and Carbon Monoxide, 898
Exhaust emissions from four-stroke lawn mower engines, 945
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Application of computational methods for the determination of traffic emissions, 1204
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Modeling
Modeling anthropogenic trends in air quality data, 66
Formulation of a second-generation relative plume visibility model, 176
Preceptor model applied to patterns in space (RMAPS) Part I-Model description, 216
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524
Comparing neural networks and regression models for ozone forecasting, 653
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189
Application of computational methods for the determination of traffic emissions, 1204
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
São Paulo aerosol characterization study, 1297

Moisture Balance
Evaluating design options for biofilters, 37

Monitoring
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194

Comparison of selenium and sulfur at remote sites throughout the United States, 204
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Redundant measurements of urban air monitoring networks in air quality reporting, 614
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Morphology
Evaluation of dust generated from basic oxygen furnaces steel making, 716

Motor Vehicles
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
An analysis of Michigan and California CO remote sensing measurements, 601
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Multimedia
Simultaneous treatment of industrial wastewater and vent streams, 961

Municipal Collection
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061

Municipal Solid Waste
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Mutagenicity
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

NAAQS
Attainment flip-flops and the achievability of the ozone air quality standard, 620
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

National Parks
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Natural Emissions
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

New Jersey
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592

Nitrate
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Nitric Oxide
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

Nitrogen Dioxide
Characterization of air quality problems in five finnish indoor ice arenas, 1979
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Nitrogen Oxides
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Formation and transport of the Madrid ozone plume, 766
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Nitrous Oxide
Stabilization of the greenhouse impact caused by anthropogenic emissions from Nordic Countries, 806

NMOC
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Nocturnal Drainage Flows
Summer transport patterns affecting the Mohave power project emissions, 383

Nonattainment
Attainment flip-flops and the achievability of the ozone air quality standard, 620

Non-Methane Hydrocarbons
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030

Non-Methane Organic Gases
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Non-Road Sources
Exhaust emissions from four-stroke lawn mower engines, 945

Non-Traditional Sources
Exhaust emissions from four-stroke lawn mower engines, 945

Odor Control
Bioconversion of methylamine in biofilters, 58

Odors
Evaluating design options for biofilters, 37

Opacity
Formulation of a second-generation relative plume visibility model, 176

Organic Aerosol
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Organic Carbon Aerosol
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212

Oxygen Combustion System
Combustion and gasification properties of plastics particles, 861

Oxygen Sensing
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

Ozone
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Attainment flip-flops and the achievability of the ozone air quality standard, 620
Comparing neural networks and regression models for ozone forecasting, 653
Measurement of ozone concentrations in ambient air using a badge-type passive monitor, 697
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030

Ozone Boundary Layer
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030

Ozone Dry Deposition
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in Southern Europe, 792

Ozone Formation
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon Sensitivity: Case studies from Atlanta, New York, and Los Angeles, 1030

Ozone, Tropospheric
Attainment flip-flops and the achievability of the ozone air quality standard, 620
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Passive Samplers
Measurement of ozone concentrations in ambient air using a badge-type passive monitor, 697
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

PCBs
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Percent Removal
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976

Perfluorocarbon Tracers
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990

Personal Monitoring
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Pesticides
Biotransformation of herbicides in the presence of various electron acceptors, 690

Phillips Number
Simultaneous treatment of industrial wastewater and vent streams, 961

Photochemical Pollutants
Formation and transport of the Madrid ozone plume, 766

Pine Barks
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

PIXE
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Plasma Systems
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Plastics
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Plumes
Formulation of a second-generation relative plume visibility model, 176

$PM_{2.5}$
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

PM_{10}
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
An assessment of source contributions to ambient aerosols in central Taiwan, 501
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784
Airborne hexavalent chromium in Southwestern Ontario, 905
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238
São Paulo aerosol characterization study, 1297

Point Source Emissions
A new method for measuring vapor emissions, 1161

Polar Organic Compounds
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Policy Decision Making
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976

Pollutant Concentration
Modeling anthropogenic trends in air quality data, 66
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Redundant measurements of urban air monitoring networks in air quality reporting, 614
Formation and transport of the Madrid ozone plume, 766
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Pollution Control Equipment
Status and future of baghouses in the utility industry, 704

Pollution Prevention
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
Integrated environmental control modeling of coal-fired power systems, 1180

Polycyclic Aromatic Hydrocarbons
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953

Polyethylene
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Population Exposure
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Power Plants
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269
Summer transport patterns affecting the Mohave power project emissions, 383
Combustion and gasification properties of plastics particles, 861

Primary Ozone Standard
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Principal Organic
Air quality monitoring during construction and initial occupation of a new building, 890

Processing
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Project MOHAVE
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Public Health
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Health risk assessment of fluctuating concentrations using lognormal models, 1152

Public Owned Treatment Works
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Receptor Modeling
Preceptor model applied to patterns in space (RMAPS) Part I-Model description, 216
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Preceptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Receptor Models
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Recycling
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061

Regulatory Policy
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468

Remediation
Biotransformation of herbicides in the presence of various electron acceptors, 690
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189

Remote Sensing
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582
An analysis of Michigan and California CO remote sensing measurements, 601

Research and Development
Methods for real-time, in situ measurement of aerosol light absorption, 157

Residences
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Residence Time
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Respiratory Health
Airborne hexavalent chromium in Southwestern Ontario, 905

Resurfacer Emission Control
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

Roadways
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

Rotary Kiln
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189

Rural Sites
Spatial and temporal patterns in particle data measured during the MOHAVE Study, 119
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
Comparison of selenium and sulfur at remote sites throughout the United States, 204
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411

Samplers, Mobile
Formation and transport of the Madrid ozone plume, 766

Sampling
Analyses of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136
Fine particulate organic material at Meadview during the project MOHAVE

summer intensive study, 357
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682
Air quality monitoring during construction and initial occupation of a new building, 890
A new method for measuring vapor emissions, 1161

Sampling Artifacts
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Scattering
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Scrubbers
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Seasonal Effects
Spatial and temporal patterns in particle data measured during the MOHAVE Study, 119
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
Comparison of selenium and sulfur at remote sites throughout the United States, 204
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

SITE Program Demonstration
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Size Distribution
Analyses of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136

Small Engines
Exhaust emissions from four-stroke lawn mower engines, 945

Smog
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

SO_2 Absorption
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Software Products
Integrated environmental control modeling of coal-fired power systems, 1180

Soil
Partitioning of arsenic species in fine-grained soils, 481
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Soil Dust
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Solid Waste
Combustion and gasification properties of plastics particles, 861
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Solvent Extraction
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Soot
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317

Sorbent Tube
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20

Source Apportionment
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Source Profiles
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Spatial Variability
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517

Statistical Analysis
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517

Statistics
Calculating removal efficiencies: Confidence limits and hypothesis tests, 976

Sulfate
Preceptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from project MOHAVE, 220
Preceptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226
Characterization of the southwestern desert aerosol, Meadview, AZ, 344
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Sulfur
Analyses of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136

Sulfur Dioxide
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331

Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

Summer Monsoonal Patterns
Summer transport patterns affecting the Mohave power project emissions, 383

Superfund
On the impact of future land use assumptions on risk analysis for superfund sites, 881
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Superfund Innovative Technology Evaluation (SITE)
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Synoptic Classification Scheme
Summer transport patterns affecting the Mohave power project emissions, 383

Technology Development
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Tertiary Butyl Alcohol (TBA)
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Test Method
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Testing
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

Thermal Desorption
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20

Thermal Treatment
Combustion and gasification properties of plastics particles, 861
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189
Destruction of chlorofluorocarbons in a cement kiln, 1220

Thin-Film Evaporation
Simultaneous treatment of industrial wastewater and vent streams, 961

Throughfall Rates
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Toluene
Screening of biofiltering material for VOC treatment, 674

Toxic Air Pollutants
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Airborne hexavalent chromium in Southwestern Ontario, 905

Toxic Gases
Henry's law coefficient of hydrazoic acid, 1216

Toxic Metals
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
Airborne hexavalent chromium in Southwestern Ontario, 905
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Toxics
Biotransformation of herbicides in the presence of various electron acceptors, 690

Toxic Wastes
Henry's law coefficient of hydrazoic acid, 1216
Toxicity and Mutagenicity of Component Classes of Oils Isolated from Soils at Petroleum-and Creosote-Contaminated Sites, 1250

Tracers
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Traffic
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491
Application of computational methods for the determination of traffic emissions, 1204

Trajectory Analysis
Clean air corridors: A geographic and meteorologic characterization, 403

Treatment Technologies
Biotransformation of herbicides in the presence of various electron acceptors, 690
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Trend Analysis
Modeling anthropogenic trends in air quality data, 66
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

Urban Areas
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in Soil, 592
Redundant measurements of urban air monitoring networks in air quality reporting, 614
Comparing neural networks and regression models for ozone forecasting, 653
Formation and transport of the Madrid ozone plume, 766
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

Urine
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

U.S.EPA
Analysis of PM_{10} trends in the United States from 1988 through 1995, 1070

Validation of Canopy 1-D models
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in Southern Europe, 792

Vapors
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Venturi Scrubber
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Visibility
Emission estimates for assessing visual air quality on the Colorado Plateau, 185
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395
Clean air corridors: A geographic and meteorologic characterization, 403

Visibility Impairment
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212

Volatile Organic Compounds (VOCs)
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20
Evaluating design options for biofilters, 37
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608
Kinetic behavior of the toluene biofiltration process, 664
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710
Formation and transport of the Madrid ozone plume, 766
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871
Air quality monitoring during construction and initial occupation of a new building, 890
Water content dynamics in biofiltration: The role of humidity and microbial heat generation, 936
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111
Transient behavior of the biofilters for toluene removal, 1142
A new method for measuring vapor emissions, 1161
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

VOC Control
Screening of biofiltering material for VOC treatment, 674

Waste Disposal
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49
Combustion and gasification properties of plastics particles, 861

Waste Management
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103

Waste Reduction
Combustion and gasification properties of plastics particles, 861
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Waste Site Operations
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Waste Treatment
Biotransformation of herbicides in the presence of various electron acceptors, 690
Combustion and gasification properties of plastics particles, 861
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Wastewater Treatment
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Wind Conditions
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

White Phosphorus
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103

X-ray Diffraction
Evaluation of dust generated from basic oxygen furnaces steel making, 716

XRF
Ambient air concentrations of fine (PM2.5) manganese in U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Author Index

Adams, K.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Alba, N.
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Allen, G.
Characterization of the southwestern desert aerosol, Meadview, AZ, 344
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Alm, S.
Characterization of air quality problems in five finnish indoor ice arenas, 1079

Almbauer, R.
Application of computational methods for the determination of traffic emissions, 1204

Alonso, C.D.
São Paulo aerosol characterization study, 1297

Aneja, V.P.
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

Anh, V.
Modeling anthropogenic trends in air quality data, 66

Anshumali
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Apam, D.
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Arnott, W.P.
Methods for real-time, in situ measurement of aerosol light absorption, 157

Artiñano, B.
Formation and transport of the Madrid ozone plume, 766

Arunachalam, S.
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Ashley, D.
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Azzi, M.
Modeling anthropogenic trends in air quality data, 66

Balentine, H.W.
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Barbiaux, M.
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Baldasano, J.M.
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Barlow, A.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Belian, T.C.
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426

Bell, R.W.
Airborne hexavalent chromium in southwestern Ontario, 905

Bergstrom, R.W.
Formulation of a second-generation reactive plume and visibility model, 176

Berkenpas, M.B.
Integrated environmental control modeling of coal-fired power systems, 1180

Betterton, E.A.
Henry's law coefficient of hydrazoic acid, 1216

Bishop, P.L.
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Blanchard, C.L.
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Braathen, O.A.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Brauer, M.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Broholm, K.
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Brook, J.R.
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2

Brothers, H.S.
An alternate compliance strategy for Title III of the 1990 Clean Air Act amendments, 468

Buckley, T.J.
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Burnett, R.T.
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Byrne, S.V.
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Cadle, S.H.
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426

Cahill, T.A.
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194

Caka, F.M.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Cana, L.
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

JOURNAL
OF THE AIR & WASTE MANAGEMENT ASSOCIATION VOLUME 47

Cardelino, C.
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon sensitivity: case studies from Atlanta, New York, and Los Angeles, 1030

Chan, C.-C.
Redundant measurements of urban air monitoring networks in air quality reporting, 614

Chattopadhyay, G.
Evaluation of dust generated from basic oxygen furnace steel making, 716

Chestnut, L.G.
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395

Chen, W.-C.
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Cheng, W.-H.
Screening of biofiltering material for VOC treatment, 674

Cherry, R.S.
Water content dynamics in biofiltration: the role of humidity and microbial heat generation, 936

Chock, D.P.
Attainment flip-flops and the achievability of the ozone air quality standard, 620

Chou, M.-S.
Bioconversion of methylamine in biofilters, 58
Screening of biofiltering material for VOC treatment, 674

Chow, J.C.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Chyang, C.-S.
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49

Cicero-Fernández, P.
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898

Comrie, A.C.
Comparing neural networks and regression models of ozone forecasting, 653

Constantinou, E.
Formulation of a second-generation reactive plume and visibility model, 176

Cui, W.
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Dalager, A.
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Dakins, M.
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Dann, T.F.
The relationship among TSP, PM_{10}, $PM_{2.5}$, and inorganic constituents of atmospheric particulate matter at multiple Canadian locations, 2
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Darlington, T.L.
Analysis of PM_{10} trends in the Unites States from 1988 through 1995, 1070

Day, D.E.
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411

Deaton, M.L.
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Dennis, R.L.
Economic benefits of improvements in visibility: Acid rain provisions of the 1990 Clean Air Act amendments, 395

Deshusses, M.A.
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Dickson, R.J.
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Dolislager, L.J.
The effect of California's wintertime oxygenated fuels program on ambient carbon monoxide concentrations, 775

Drahonovska, H.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Drees, L.
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Du, A.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101

Duc, H.
Modeling anthropogenic trends in air quality data, 66

Dunne, I.J.
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Eatough, D.J
Regional source profiles of sources of SO_x at the Grand Canyon during project MOHAVE, 101
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Eatough, M.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Eatough, N.L.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Fine particulate organic material at Meadview during the Project MOHAVE summer intensive study, 357
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Eklund T.
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

Eldred, R.A.
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194
Comparison of selenium and sulfur at remote sites throughout the United States, 204

Everett, J.W.
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061

Farber, R.J.
Spatial and temporal patterns in particle data measured during the MOHAVE study, 119
Summer transport patterns affecting the Mohave power project emissions, 383

Farouk, B.
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189

Fathikalajahi, J.
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Finley, B.L.
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in soil, 592
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Flocchini, R.G.
Composition of $PM_{2.5}$ and PM_{10} aerosols in the IMPROVE Network, 194

Frankenhaeuser, M.
Combustion and gasification properties of plastics particles, 861

Frey, H.C.
Integrated environmental control modeling of coal-fired power systems, 1180

Gabele, P.
Exhaust emissions from four-stroke lawn mower engines, 945

García, I.
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Gasso, S.
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Gebhart, K.A.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
Clean air corridors: A geographic and meteorologic characterization, 403

Georgopoulos, P.G.
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Giles, M.
An analysis of Michigan and California CO remote sensing measurements, 601

Gillespie, P.
Formulation of a second-generation reactive plume and visibility model, 176

Gimeno, L
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

Godinho, R.
São Paulo aerosol characterization study, 1297

Goldstein, R.
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Gorse, R.A.
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426
An analysis of Michigan and California CO remote sensing measurements, 601

Gostomski, P.A.
Water content dynamics in biofiltration: the role of humidity and microbial heat generation, 936

Graboski, M.S.
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

Green, M.C.
Analysis of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136
Summer transport patterns affecting the Mohave power project emissions, 383
Clean air corridors: A geographic and meteorologic characterization, 403

Guise-Bagley, L.
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Hwang, S.J.
Transient behavior of the biofilters for toluene removal, 1142

Harris, M.A.
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in soil, 592

He, D.
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon sensitivity: case studies from Atlanta, New York, and Los Angeles, 1030

Hedley, M.
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Henry, R.C.
Receptor model applied to patterns in space (RMAPS) Part I-Model description, 216
Receptor model applied to patterns in space (RMAPS) Part II-Apportionment of airborne particulate sulfur from Project MOHAVE, 220
Receptor model applied to patterns in space (RMAPS) Part III-Apportionment of airborne particulate sulfur in western Washington State, 226

Hernández, E.
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

Heuss, J.M.
Analysis of PM_{10} trends in the Unites States from 1988 through 1995, 1070

Hildemann, L.
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Hindawi, M.A.
Simultaneous treatment of industrial wastewater and vent streams, 961

Hipfner, J.C.
Airborne hexavalent chromium in southwestern Ontario, 905

Ho, J.-A.
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49

Hoff, R.
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Hoffer, T.E.
Summer transport patterns affecting the Mohave power project emissions, 383

Hoffman, D.
An analysis of Michigan and California CO remote sensing measurements, 601

Holbrook, B.D.
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

Holcombe, J.T.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Holdren, M.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Howe, G.B.
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Huber, A.H.
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Huhn, G.
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Hupa, M.
Combustion and gasification properties of plastics particles, 861

Husar, R.B.
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331

Hwang, J.-S.
Redundant measurements of urban air monitoring networks in air quality reporting, 614

Hwang, S.-J.
Kinetic behavior of the toluene biofiltration process, 664

Imhoff, R.E.
The use of photochemical indicators to evaluate Ozone-NO_x-Hydrocarbon sensitivity: case studies from Atlanta, New York, and Los Angeles, 1030

Iwasaki, Y.
Destruction of chlorofluorocarbons in a cement kiln, 1220

Iyer, H.K.
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370

Jaakkola, PT.
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

James, R.
An analysis of Michigan and California CO remote sensing measurements, 601

Jantunen, M.J.
Characterization of air quality problems in five finnish indoor ice arenas, 1079

Jayanty, R.K.M.
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Jefcoat, I.A.
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103

Jensen, D.T.
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Joseph, J.M.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Kahl, J.D.W.
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Kahlbaum, D.F.
Analysis of PM_{10} trends in the Unites States from 1988 through 1995, 1070

Kalagnanam, J.R.
Integrated environmental control modeling of coal-fired power systems, 1180

Karamchandani, P.
Formulation of a second-generation reactive plume and visibility model, 176
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Karlsson, M.
Combustion and gasification properties of plastics particles, 861

Kastenberg, W.E.
On the impact of future land use assumptions on risk analysis for superfund sites, 881

Katsumata, P.T.
On the impact of future land use assumptions on risk analysis for superfund sites, 881

Keener, T.C.
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Kenefick, S.L.
Toxicity and mutagenicity of component classes of oils isolated from soils at petroleum-and creosote-contaminated sites, 1250

Khang, S.J.
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Kjeldsen, P.
Attenuation of methane and nonmethane organic compounds in landfill gas affected soils, 1268

Kjoergaard, S.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Knaebel, K.S.
A new method for measuring vapor emissions, 1161

Koutrakis, P.
Characterization of the southwestern desert aerosol, Meadview, AZ, 344
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Kreidenweis, S.M.
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411

Ku, J.Y.
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Kuhlmeier, P.D.
Partitioning of aresenic species in fine-grained soils, 481

Kuttruff, C.E.
Simultaneous treatment of industrial wastewater and vent streams, 961

Lacorte, T.
Characterization of municipal solid waste incineration residues from facilities with different air pollution control systems, 1170

Lagomarsino, R.J.
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990

Larjava, K.T.
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Latner, N.
A tethered balloon air sampling system for determining the vertical distribution of atmospheric tracers, 990

Laulainen, N.
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212

Law, P.L.
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Lawson, D.R.
Real-world vehicle emissions: A summary of the sixth coordinating research council on-road vehicle emissions workshop, 426

Lee, K.
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Leson, G.
Evaluating design options for biofilters, 37

Levine, S.P.
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Lewis, L.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357
Precision and accuracy in the determination of sulfur oxides, fluoride, and spherical aluminosilicate fly ash particles in Project MOHAVE, 455

Liebergeld, G.
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Lioy, P.J.
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Lipfert, F.W.
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Liu, D.
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Long, J.R.
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898

López, A.
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in southern Europe, 792

Loranger, S.
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

Lurmann, F.W.
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Machir, J.
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167
Fine particulate organic material at Meadview during the project MOHAVE summer intensive study, 357

Macias, E.S.
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Malm, W.C.
Spatial and temporal patterns in particle data measured during the MOHAVE study, 119
Source apportionment of sulfur and light extinction using receptor modeling techniques, 250
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370
Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks, 411

Mangelson, N.F.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Marrack, D.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Martins, M.H.R.B.
São Paulo aerosol characterization study, 1297

Mathews, T.B.
Isothermal oxidation of white phosphorus dispersed in water in a stirred-tank reactor, 1103

McAlinden, K.
An analysis of Michigan and California CO remote sensing measurements, 601

McCormick, R.L.
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

McCurdy, T.R.
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

McLaren, R.
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

McMurry, P.
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Meckes, M.C.
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Merritt, R.L.
Status and future of baghouses in the utility industry, 704

Michelsen, R.
Evaluating design options for biofilters, 37

Middleton, P.
DAQM-simulated spatial and temporal differences among visibility, PM, and other air quality concerns under realistic emission change scenarios, 302

Mihalikova, E.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Miskovic, P.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Moosmüller, H.
Methods for real-time, in situ measurement of aerosol light absorption, 157

Moss, P.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Neff, W.D.
The Denver Brown Cloud Studies from the perspective of model assessment needs and the role of meteorology, 269

Newlin, A.W.
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

Niehus, B.
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Nozaki, A.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Nylund, N.O.
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

Oliver, W.R.
Emission estimates for assessing visual air quality on the Colorado Plateau, 185

Orton, T.L.
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Pai, P.
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Park, D.Y.
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Parker, E.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Parker, W.J.
A multi-parameter sensitivity analysis of a model describing the fate of volatile organic compounds in trickling filters, 871

Pasanen, P.
Characterization of air quality problems in five finnish indoor ice arenas, 1079

Pennanen, A.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Pennanen, A.S.
Characterization of air quality problems in five finnish indoor ice arenas, 1079
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087

Pezda, S.
Attainment flip-flops and the achievability of the ozone air quality standard, 620

Phillips, J.B.
Simultaneous treatment of industrial wastewater and vent streams, 961

Pitchford, M.
Analysis of sulfur aerosol size distributions for a forty-day period in summer 1992 at Meadview, Arizona, 136

Plaza, J.
Formation and transport of the Madrid ozone plume, 766

Poirot, R.L.
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Pollard, S.J.
Toxicity and mutagenicity of component classes of oils isolated from soils at petroleum-and creosote-contaminated sites, 1250

Porter, P.S.
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Pradhan, S.P.
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710

Prah, J.D.
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739

Pryor, S.C.
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Pucher, K.
Application of computational methods for the determination of traffic emissions, 1204

Pujadas, M.
Formation and transport of the Madrid ozone plume, 766

Qing-Xiang, Z.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Rabbe, D.E.
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in soil, 592

Rao, S.T.
Small sample properties of nonparametric bootstrap t confidence intervals, 1197

Ray, A.K.
Evaluation of dust generated from basic oxygen furnace steel making, 716

Ray, S.K.
Evaluation of dust generated from basic oxygen furnace steel making, 716

Rees, L.B.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101

The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Reiss, R.
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Rickman, Jr., E.E.
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Riley, P.
Curbside collection of recyclable material: Simulation of collection activities and estimation of vehicle and labor needs, 1061

Robarge, W.P.
Nitrogen oxide flux from an agricultural soil during winter fallow in the upper coastal plain of North Carolina, U.S.A., 800

Roberts, P.T.
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Robinson, J.L.
Henry's law coefficient of hydrazoic acid, 1216

Rodríguez, R.
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in southern Europe, 792

Rogers, C.F.
Methods for real-time, in situ measurement of aerosol light absorption, 157

Romano, J.
São Paulo aerosol characterization study, 1297

Romieu, I.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Roos, A.A.
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Ross, D.M.
Human visual sensitivity to plumes with a gaussian luminance distribution: Experiments to develop and empirical probability of detection model, 370

Ross, J.D.
Effect of humidity on heavy-duty transient emissions from diesel and natural gas engines at high altitude, 784

Rubin, E.S.
Integrated environmental control modeling of coal-fired power systems, 1180

Rui-Jin, S.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Ruíz, M.E.
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Sakiyama, S.
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Salonen, R.O.
Characterization of air quality problems in five finnish indoor ice arenas, 1079
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Saltzman, B.E.
Health risk assessment of fluctuating concentrations using lognormal models, 1152

Sánchez, M.
Time trends and seasonal variation of the rainwater chemical composition in Spain, 811

Sánchez, M.L.
Ozone dry deposition in a semi-arid steppe and in a coniferous forest in southern Europe, 792

Savolainen, I.
Stabilization of the greenhouse impact caused by anthropogenic emission from Nordic countries, 806

Saxena, P.
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Saylor, E.
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Schichtel, B.A.
Regional simulation of atmospheric pollutants with the CAPITA Monte Carlo model, 331

Schroer, T.
Development of emission factors for ethylene-vinyl acetate and ethylene-methyl acrylate copolymer processing, 1111

Schulz, H.
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Schulze, R.H.
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Schüürmann, G.
Determination of throughfall rates on the basis of pine bark loads: Results of a pilot field study, 510

Scott, P.K.
Background air concentrations of Cr(VI) in Hudson County, New Jersey: Implications for setting health-based standards for Cr(VI) in soil, 592
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Seigneur, C.
Formulation of a second-generation reactive plume and visibility model, 176

Shang, S.-H.
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49

Sheth, A.
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Shiu, W.-Z.
Bioconversion of methylamine in biofilters, 58

Sillman, S.
The use of photochemical indicators to evaluate ozone-NO_x-hydrocarbon sensitivity: case studies from Atlanta, New York, and Los Angeles, 1030

Simpson, R.
Visibility and aerosol composition in the Fraser Valley during REVEAL, 147

Singleton, D.L.
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Sinisalo, J.
Stabilization of the greenhouse impact caused by anthropogenic emission from Nordic countries, 806

JOURNAL
OF THE AIR & WASTE MANAGEMENT ASSOCIATION VOLUME 47

Sioutas, C.
Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas, 682

Sissler, J.F.
Characteristics of winter and summer aerosol mass and light extinction on the Colorado Plateau, 317

Sisson, J.B.
Water content dynamics in biofiltration: the role of humidity and microbial heat generation, 936

Sloane, C.
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Slonecker, T.
Ambient air concentrations of fine ($PM_{2.5}$) manganese in the U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Smith, S.
Measurement of ozone concentrations in ambient air using a badge-type passive monitor, 697

Soufuki, M.
Destruction of chlorofluorocarbons in a cement kiln, 1220

Spengler, J.D.
Improvement of air quality in a small indoor ice arena by effective emission control in ice resurfacers, 1087
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Srivastava, V.J.
A pilot-scale demonstration of an innovative soil remediation process: Air emissions quality, 710

Stephens, R.D.
An analysis of Michigan and California CO remote sensing measurements, 601

Sturm, P.J.
Application of computational methods for the determination of traffic emissions, 1204

Sudy, C.
Application of computational methods for the determination of traffic emissions, 1204

Suh, H.H.
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Sun, B.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101

Sung, H.-M.
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Sykes, I.
Formulation of a second-generation reactive plume and visibility model, 176

Taheri, M.
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Talaie, M.R.
Mathematical modeling of SO_2 absorption in a venturi scrubber, 1211

Tang, H.-M.
Kinetic behavior of the toluene biofiltration process, 664
Transient behavior of the biofilters for toluene removal, 1142

Tatsuichi, S.
Destruction of chlorofluorocarbons in a cement kiln, 1220

Teng, H.
Characterization of waste tire incineration in a prototype vortexing fluidized bed combustor, 49

Thompson, R.J.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Tillman, J.
Removal of PCBs from a contaminated soil using CF-Systems, solvent extraction process, 1119

Tormonen, K.E.
Field measurements of flue gases from combustion of miscellaneous fuels using a low-resolution FTIR gas analyzer, 1284

Trexler, E.
Assessing the relative contribution of biogenic and fossil processes to visibility-scattering aerosols found in remote areas: Near-term organic research program, 212

Tsuzuki, T.
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Turner, D.B.
Identification of an accurate soil suspension/dispersion modeling method for use in estimating health-based soil cleanup levels of hexavalent chromium in chromite ore processing residues, 753

Turpin, B.J.
Characterization of the southwestern desert aerosol, Meadview, AZ, 344

Ueno, H.
Destruction of chlorofluorocarbons in a cement kiln, 1220

Valicenti, J.A.
Air quality monitoring during construction and initial occupation of a new building, 890

VanderGheynst, G.B.
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

VanderGheynst, J.S.
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

Van Lith, C.
Evaluating design options for biofilters, 37

Vann Bush, P.
Status and future of baghouses in the utility industry, 704

Vasconcelos, L.
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Vedal, S.
Critical review: Ambient particles and health: Lines that divide, 551
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Vega, E.
Application of a chemical mass balance receptor model to respirable particulate matter in Mexico City, 524

Venkatram, A.
Formulation of a second-generation reactive plume and visibility model, 176
The development of a model to examine source-receptor relationships for visibility on the Colorado Plateau, 286

Vostal, J.J.
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Walker, L.P.
Development and analysis of oxygen-sensing probes for in situ monitoring of unsaturated solid-state biodegradation processes, 1041

Wallace, J.
Calculating removal efficiencies: confidence limits and hypothesis tests, 976

Wallace, L.
Ambient air concentrations of fine ($PM_{2.5}$) manganese in the U.S. national parks and in California and Canadian cities: The possible impact of adding MMT to unleaded gasoline, 642

Walsh, P.A.
Summer transport patterns affecting the Mohave power project emissions, 383

Wang, C.-S.
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Wang, D.
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Wang, G.
Biotransformation of herbicides in the presence of various electron acceptors, 690

Wang, S.
Alternative metrics for assessing the relative effectiveness of No_x and VOC emission reductions in controlling ground-level ozone, 838

Watson, J.G.
Regional source profiles of sources of SO_x at the Grand Canyon during Project MOHAVE, 101
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Wei, C.-C.
An assessment of source contributions to ambient aerosols in central Taiwan, 501

Wei, Y.L.
PAH Emissions from the fluidized-bed incineration of an industrial sludge, 953

Weimin, Jiang
Comparison of organic compound compositions in the emissions inventory and ambient data for the Lower Fraser Valley, 851

Wenger, J.
Air quality monitoring during construction and initial occupation of a new building, 890

White, W.H.
The relationship between atmospheric transport and the particle scattering coefficient at the Grand Canyon, 419

Whitfield, J.K.
Interlaboratory study of a test method measuring total volatile organic compound content of consumer products, 608

Wilber, G.G.
Biotransformation of herbicides in the presence of various electron acceptors, 690

Wilkerson, T.
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Williams, N.W.
The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, 167

Wilson, W.E.
Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies, 1238

Winebrake, J.J.
A comparative analysis of emissions deterioration for in-use alternative fuel vehicles, 1291

Winer, A.M.
Effects of grades and other loads on on-road emissions of hydrocarbons and carbon monoxide, 898

Winkleman, B.C.
Destruction of low levels of volatile organic compounds in dry air streams by an electron-beam generated plasma, 1276

Winkler, S.L.
Attainment flip-flops and the achievability of the ozone air quality standard, 620

Wolff, G.T.
Analysis of PM_{10} trends in the Unites States from 1988 through 1995, 1070

Woolfenden, E.
Monitoring VOCs in air using sorbent tubes followed by thermal desorption-capillary GC analysis: Summary of data and practical guidelines, 20

Wu, C.H.
PAH emissions from the fluidized-bed incineration of an industrial sludge, 953

Wu, X.A.
Formulation of a second-generation reactive plume and visibility model, 176

Wyzga, R.E.
Air pollution and mortality: The implications of uncertainties in regression modeling and exposure measurement, 517
1997 Critical review discussion: Ambient particles and health: Lines that divide, 995

Xu, Y
Nitrogen dioxide in indoor ice skating facilities: An international survey, 1095

Yang, L.
Modeling of solid particle flow and heat transfer in rotary kiln calciners, 1189

Yost, M.G.
Evaluation of virtual source beam configurations for rapid tomographic reconstruction of gas and vapor concentrations in workplaces, 582

Zayed, J.
Environmental contamination and human exposure to airborne total and respirable manganese in Montreal, 983

Zelenka, M.P.
Evaluation of a probabilistic exposure model applied to carbon monoxide (pNEM/CO) using Denver personal exposure monitoring data, 491

Zemanek, M.G.
Toxicity and mutagenicity of component classes of oils isolated from soils at petroleum-and creosote-contaminated sites, 1250

Zevehoven, R.
Combustion and gasification properties of plastics particles, 861

Zhou, J.
Measurement of ozone concentrations in ambient air using a badge-type passive monitor, 697

Zhu, H.
The effectiveness of aeration recirculation in controlling VOC emissions from publicly owned treatment works, 1259

Ziman, S.D.
The application of exposure-based criteria in developing alternative primary ambient ozone standards, 1051

Zuber, L.
Comparative scale-up and cost estimation of a biological trickling filter and three-phase airlift bioreactor for the removal of methylene chloride from polluted air, 969

Zweidinger, R.A.
Body burden measurements and models to assess inhalation exposure to Methyl Tertiary Butyl Ether (MTBE), 739